



GUARANTEED ANALYSIS

Total Organic Matter	: 20 % (w/v)
Total Nitrogen (N)	: 6 % (w/v)
Urea Nitrogen (NH ₂ -N)	: 6 % (w/v)
Water Soluble Potassium Oxide (K ₂ O)	: 6 % (w/v)
Water Soluble Calcium Oxide (CaO)	: 2,5 % (w/v)
Water Soluble Boron (B)	: 0,01 % (w/v)
Water Soluble Copper (Cu)	: 0,002 % (w/v)
Water Soluble Iron (Fe)	: 0,02 % (w/v)
Water Soluble Manganese (Mn)	: 0,01 % (w/v)
Water Soluble Molybdenum (Mo)	: 0,05 % (w/v)
Water Soluble Zinc (Zn)	: 0,002 % (w/v)
Free Amino Acid	: 2 % (w/v)

Liquid Organomineral Fertiliser With Nitrogen And Potassium

MULTI-X

Liquid Macro-Micro
Nutrient Fertilisers

FEATURES

MULTI-X is a special organomineral fertiliser developed to provide organic material needs of the plant and the soil, as well as macro and micro elements. Its organamineral blend helps with plant growth, as well as increasing resistance to climate conditions like heat or cold, and increasing fruit yield. MULTI-X can be used on many vegetables and fruits, both in greenhouses and on open fields.

In addition, the high level of easily absorbable and high quality free amino acid content of MULTI-X provides a noticeable difference in plant growth, ripening and fruit yield.

BENEFITS

MULTI-X is a special product rich in Amino Acids, derived from wet plants. In perennial plants, it is suitable to start to use with plant sap water movement. It can be safely used in annual plants, from planting seedlings or from plants bearing 3 - 5 leaves.

MULTI-X stimulates the growth tip of the plant due to its special content.

MULTI-X increases overground plant shoot activity and re-sprouting. It can be safely used in plants where growth stagnation is observed.

MULTI-X increases the resistance of plants against adverse environmental conditions (soil salinity, heat, cold, aquatic plant, etc.).

MULTI-X increases the storage, quality and shelf life of fruits.

MULTI-X is used safely in combination with phytosanitary products, fertilisers and plant growth regulators.

MULTI-X provides an indispensable solution for fruit growth and colour problems in its ideal formulation ratios.

MULTI-X promotes root development and increases fruit formation.

MULTI-X increases fruit growth in cold periods, and prevents choking especially in pepper.

MULTI-X helps transfusion of synthesized organic materials to other plant organs, where they are needed in its potassium content.

MULTI-X facilitates plant growth into the ripe stage.

MULTI-X prevents fruit gaps. Promotes formation of full and regular fruits, thus increasing market value of the crops.

USAGE AREA, FORM, TIME AND QUANTITY

USAGE AREA	APPLICATION TIME	SOIL APPLICATION	FOLIAR APPLICATION
All Greenhouses and All Open Field and Winter Vegetables	1 - 2 applications with drip irrigation 10 days after planting. From pre-flowering until the harvest, it can be applied with foliar application at regular intervals.	20 Lt. / Ha	250 - 300 cc. / 100 Lt. Water
Strawberry and Ornamental Plants	Begins with vegetative growth and apply at 10 - 15 day intervals throughout the season.	20 Lt. / Ha	250 - 300 cc. / 100 Lt. Water
Vineyard	From the beginning of the shoot until the harvesting period, it can be applied by drip irrigation when needed. It can also be applied by foliar application at regular intervals until fruit set period.	20 Lt. / Ha	250 - 300 cc. / 100 Lt. Water
Corn and Sunflower	Every 7 - 10 days from bud break to bloom. Use while Nitrogen fertilization is necessary.	20 Lt. / Ha	250 - 300 cc. / 100 Lt. Water
Cereals and Forage Crops	During the tillering period in early spring.	20 Lt. / Ha	300 - 350 cc. / 100 Lt. Water
All Stone and Pome Fruit Trees and Nut Crops	From beginning of blooming until the harvesting, it can be applied at regular intervals.	20 Lt. / Ha	150 - 200 cc. / 100 Lt. Water
Cotton	It can be used safely in every period needed from when 8 - 10 leaves appear.	25 Lt. / Ha	500 cc. / 100 Lt. Water

MIXABILITY

It is recommended to conduct small container compatibility tests before mixing with the other products. To be used only if required, and do not exceed the recommended dosages.

PACKAGING:

