



GUARANTEED ANALYSIS

Organic Matter	: 15 % (w/w)
Total Nitrogen (N)	: 5 % (w/w)
Nitrate Nitrogen (NO ₃ -N)	: 5 % (w/w)
Water Soluble Potassium Oxide (K ₂ O)	: 15 % (w/w)
Water Soluble Magnesium Oxide (MgO)	: 2 % (w/w)
Water Soluble Boron (B)	: 0,03 % (w/w)
Water Soluble Copper (Cu)	: 0,03 % (w/w)
Water Soluble Iron (Fe)	: 0,04 % (w/w)
Water Soluble Manganese (Mn)	: 0,05 % (w/w)
Water Soluble Zinc (Zn)	: 0,05 % (w/w)

Liquid Organomineral Fertiliser With Nitrogen And Potassium

Liquid Macro-Micro
Nutrient Fertilisers

FEATURES

POTTASOL is a specific product that responds quickly to the Potassium (K) requirement of plants.

POTTASOL

BENEFITS

Because of its special formulation, in all stress condition (very hot, cold and salty) it can be absorbed by the plants root very easily.

When used in drip irrigation systems, the nitrate (NO₃) is negatively (-) charged in the root zone and the potassium (K) + are combined compound to form potassium nitrate (KN₃)- which are transmitted through the roots to the beams.

With foliar applications with help of the enzymes and activators, it goes through the stomas to meet the need of Potassium (K).

It directly affects the fruit growth, colour, quality and sugar ratio, for this it increases the fruit market value.

It eliminates distortion of fruit shapes and helps standardize the fruit skin.

There is no consequences with overuse. Over dose is suitable for sudden growth. There is no means of burning.

USAGE AREA, FORM, TIME AND QUANTITY

USAGE AREA	APPLICATION TIME	SOIL APPLICATION	FOLIAR APPLICATION
Cut Flower and Strawberry	It is used throughout the season starting from planting.	15 - 20 Lt. / Ha	250 - 300 cc. / 100 Lt. Water
All Greenhouse and Open Field Vegetables	It is used throughout the season starting from planting.	20 - 25 Lt. / Ha	300 - 400 cc. / 100 Lt. Water
Melon, Watemelon and Zucchini	During seedling planting with life water and can be applied to seedling pillows.	20 - 25 Lt. / Ha	300 - 400 cc. / 100 Lt. Water
Com, Sunflower, Cotton, Soya, Chickpea, Beans, Peanut and Lentil	2 - 3 applications from 30 - 45 days prior to blooming (April-May).	20 - 25 Lt. / Ha	300 - 400 cc. / 100 Lt. Water
Vineyards	It is used throughout the season starting from sprouting.	20 Lt. / Ha	300 - 400 cc. / 100 Lt. Water
All Stone and Pome Fruit Trees	From beginning of blooming until the harvesting, it can be applied at regular intervals.	20 - 25 Lt. / Ha	250 - 300 cc. / 100 Lt. Water
Grains, Forage Plants and Green Areas	In the early spring and tillering period.	-	300 - 400 cc. / 100 Lt. Water

MIXABILITY

It is recommended to conduct small container compatibility tests before mixing with the other products. To be used only if required, and do not exceed the recommended dosages.

PACKAGING:



10 Kg



www.ce-on.com