




GUARANTEED ANALYSIS

Water Soluble Boron (B)	: 8 % (w/w)
Water Soluble Manganese (Mn) Chelated with EDTA	: 1 % (w/w)
Water Soluble Zinc (Zn) Chelated with EDTA	: 5 % (w/w)

Edta Chelated Powder Micronutrient Fertiliser

FEATURES

BOR MIX is a chemical combination of essential chelated microelements; specially indicated for optimum fertilization for an optimum nutrition, balanced and quick-response, allowing optimum crop development.

BOR MIX

BENEFITS

BOR-MIX is a mixture of liquid microelements specially prepared to prevent and eliminate boron and many micro element deficiencies in plants.

BOR-MIX is used in plants encourages the formation of stigma tube insemination.

BOR-MIX increases homogeneous fruit set when used before flowering. When used during fruit development period, fruit quality increases.

BOR-MIX is very suitable for post-harvest applications in perennial plants.

BOR-MIX increases the coloration, sales appeal and quality in the fruits thanks to its special formulation.

BOR-MIX is used safely in combination with phytosanitary products, fertilizers and plant growth regulators.

USAGE AREA, FORM, TIME AND QUANTITY

USAGE AREA	APPLICATION TIME	SOIL APPLICATION	FOLIAR APPLICATION
All Greenhouse and All Open Field Vegetables	1 week after germination. First flowering. After the first fruit formation.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
All Stone and Pome Fruit Trees	At the beginning of flowering. 15 days after fruit formation. 30 days before the harvest.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
Citrus, Banana, Fig and Olive	At the beginning of flowering. 15 days after fruit formation. 30 days before the harvest.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
Vineyards and Ornamental Plants	At the beginning of flowering. 15 days after fruit formation. 30 days before the harvest.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
Corn and Sunflower	First Application: 30 days after sowing. Second Application: as required.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
Cotton	From the beginning of first square - bud, it can be applied at regular intervals when needed.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
Industrial Plants (Soya, Peanut and Canola)	After the 3 - 5 leaf growth stage, it can be applied at regular intervals when needed.	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water
Field Crops (Wheat, Barley, Rice) and Green Areas (Golf Courses, Lawn Care and Sports Turf)	When elongation of field crops reaches 20 - 25 cm. The first period of kernel development (watery ripe and milk stages).	5 - 10 Lt. / Ha	100 - 150 Gr. / 100 Lt. Water

MIXABILITY

It is recommended to conduct small container compatibility tests before mixing with the other products. To be used only if required, and do not exceed the recommended dosages.

PACKAGING:

